

Climate-change adaptation on rangelands: linking regional exposure with diverse adaptive capacity

David D Briske^{1*}, Linda A Joyce², H Wayne Polley³, Joel R Brown⁴, Klaus Wolter⁵, Jack A Morgan⁶, Bruce A McCarl⁷, and Derek W Bailey⁸

The ecological consequences of climate change are predicted to vary greatly throughout US rangelands. Projections show warming and drying in the southern Great Plains and the Southwest, warmer and drier summers with reduced winter snowpack in the Northwest, and warmer and wetter conditions in the northern Great Plains. Primarily through their combined effects on soil water availability, these climatic changes will modify plant production and community composition, which will, in turn, affect the livelihoods of humans who rely upon livestock grazing. The ability of rangeland managers to assess risk and prepare for climate change varies greatly and reflects their different adaptive capacities. Geographically specific exposure to climate change and a diverse adaptive capacity to counteract these changes will require development of varied adaptation strategies that can accommodate the various needs and abilities of livestock managers.

Front Ecol Environ 2015; 13(5): 249–256, doi:10.1890/140266

Increasing atmospheric greenhouse-gas (GHG) concentrations have elevated global temperature by 1°C since industrialization (ca 1750) and are anticipated to result in an additional 1°C rise by mid-century, regardless of projected rates of GHG emissions (NRC 2010; IPCC 2013). Warming will be accompanied by modified amounts and patterns of precipitation, and a greater frequency and intensity of extreme weather events. At the ecosystem scale, the indirect effects of climate change – such as increased wildfire incidence and modified distributions and densities of native and invasive species, including insects and disease-causing organisms – and the direct

effects of climate change are anticipated to be similar in magnitude (NRC 2010). Collectively, the consequences of climate change, interacting with large-scale shifts in human land use, will likely modify ecosystem processes and the livelihoods that they support (Polley *et al.* 2013).

Rangelands comprise approximately 30–40% of global terrestrial area and deliver diverse ecosystem services; this land's predominant use, livestock grazing, contributes to the livelihoods of millions of humans (Sayre *et al.* 2013). These livelihoods are vulnerable to climate change from both an ecological and a socioeconomic perspective (Reynolds *et al.* 2007). Rangelands represent ecologically diverse arid and semiarid systems characterized by low plant productivity and high precipitation variability, including frequent drought. Rangeland managers often have limited financial and social capital, modest infrastructure, and few options to diversify livelihoods beyond livestock grazing, and are isolated from major urban centers and governing institutions (Sayre *et al.* 2013). The vulnerability of rangeland-based livelihoods to climate change provides a strong justification for the acceleration of planning and implementation of adaptation strategies.

Climate change is projected to have geographically diverse consequences on US rangelands (Polley *et al.* 2013). Consequently, geographically specific adaptation strategies will be required to contend with localized challenges and opportunities. The heterogeneity among individuals and groups – both within and among geographic regions – regarding the recognition, capacity, and motivation to implement various adaptation strategies represents a second critical component of climate-change adaptation (Williamson *et al.* 2012; Joyce *et al.* 2013). We contend that the interaction between the geographic specificity of climate change and the heterogeneity in the adaptive capacity

In a nutshell:

- Climate change will affect millions of people whose livelihoods are linked to livestock grazing on US rangelands
- Livestock grazing will be directly influenced by changes in the amount, nutrient content, and seasonal availability of plant production, as well as by the adverse effects of heat, water limitation, and pathogen loads on animal performance
- Geographically specific ecological consequences and diverse adaptive capacity among managers represent underappreciated and interacting components of climate-change adaptation
- Adaptation planning must recognize and accommodate both geographic specificity and varied adaptive capacity to promote effective climate-change adaptation

¹Department of Ecosystem Science and Management, Texas A&M University, College Station, TX *(dbriske@tamu.edu); ²Human Dimensions Research Program, US Department of Agriculture Forest Service, Rocky Mountain Research Station, Fort Collins, CO; ³US Department of Agriculture–Agricultural Research Service, Grassland, Soil and Water Research Laboratory, Temple, TX; continued on p 256




Figure 1. Climate-change projections for US rangelands illustrating the unique exposures to and potential consequences for livestock production in four geographic regions. Projections were generated from recent simulations of CMIP5 (courtesy of J Eischeid) and data were obtained from Taylor *et al.* (2012) and Baker and Huang (2014). Regional designations follow those of USGCRP (2009), except that the Great Plains have been divided into two separate regions, the northern and southern Great Plains, to emphasize the large differences in climate projections.

of managers – that is, their ability to respond to, create, and shape change in a system – represents an underappreciated barrier to the development and implementation of adaptation strategies. Thus, public programs, strategies, and incentives to implement climate-adaptation measures will be far more effective if they are tailored to select categories of individuals or social groups that are experiencing geographically specific climatic consequences. The objectives of this paper are to: (1) review the geographic specificity of climate change in four regions of the western US, (2) identify essential components of adaptive capacity that are relevant to rangeland managers, (3) evaluate acknowledged adaptation strategies and barriers to their implementation, and (4) highlight the geographic specificity of climate change and heterogeneous adaptive capacity among managers as interacting components of adaptation planning and implementation. Although we focus on rangelands, this conceptual framework could be applicable to other natural-resource systems, including agriculture and forestry.

■ Subcontinental climate projections and consequences

The ecological consequences of climate change are anticipated to vary greatly among geographic regions based upon current environmental conditions and unique

interactions among climate-change drivers (Figure 1; Polley *et al.* 2013). Ecosystem processes and the services that they provide will be modified by climate change and its primary drivers – elevated atmospheric carbon dioxide (CO₂) concentrations, warming, and modified precipitation regimes – on soil water availability (Knapp *et al.* 2008). Consensus projections from climate-change models for the middle of the 21st century indicate that the US southern Great Plains, the US Southwest, and northern Mexico will become warmer and drier, while the US Northwest will become warmer and drier during summer and will experience reduced snowpack in winter; at the same time, the US northern Great Plains and southern Canada will become warmer and wetter, especially during the winter (Meehl *et al.* 2007; IPCC 2013). These projections are only approximations because reliable estimates of future climates, especially for precipitation, are not yet possible at regional scales (IPCC 2013). For example, results from the recent Coupled Model Intercomparison Project Phase 5 (CMIP5) are consistent with previous temperature projections for the southwestern US (Taylor *et al.* 2012; Baker and Huang 2014; J Eischeid pers comm) but forecasted less variance for annual rainfall than previous simulations. Nevertheless, vulnerability assessments and adaptation strategies must recognize and address geographically specific exposures to effectively offset climate-change impacts (Joyce *et al.* 2013).

Southwest

In this region, warming and reduced precipitation will synergistically decrease soil water availability, thereby diminishing both the amount and nutrient content of plant production and altering plant community composition (Figure 1; Polley *et al.* 2013). Declines in water availability decrease plant growth and shorten the growing season, thus reducing available livestock forage. Collectively, changes in precipitation and temperature regimes may alter plant species composition and plant phenology to modify the seasonal availability of high-quality forage (Walther 2003; Craine *et al.* 2009). The digestibility and nutritive value of plant tissues are diminished by the higher carbon-to-nitrogen (C:N) ratio that accompanies plant growth under elevated atmospheric CO₂ conditions, and by the accelerated leaf senescence induced by water stress (Morgan *et al.* 2004). Reduced nutrient intake, higher temperatures, and more frequent heat stress are likely to further reduce livestock production and adversely affect people whose livelihoods are dependent on it (CCSP 2008; Howden *et al.* 2008). More frequent and severe droughts are anticipated to modify plant species composition and cover directly by contributing to episodic plant die-offs and indirectly through increasing the frequency of fire events. In the Mojave Desert and Great Basin over the past 20 years, increased fire frequency has converted some desert shrublands and shrub steppe communities to annual grasslands (CCSP 2008; Balch *et al.* 2013).

Southern Great Plains

Warming and drying, especially in Texas, are anticipated to reduce both plant production and nutritive content in ways similar to those described for the Southwest. These climatic changes are likely to negatively affect livelihoods by reducing stocking rates and total livestock production, and potentially by increasing production costs associated with the purchase of supplemental feed (Figure 1; Polley *et al.* 2013). Warmer temperatures may improve winter survival and increase abundance of ectoparasites, including lone star ticks (*Amblyomma americanum*) and horn flies (*Haematobia irritans*), that will further suppress livestock performance. Warming and drying are expected to modify vegetation composition and land-surface cover, including an increase in the density and cover of woody plants. Growth of both eastern redcedar (*Juniperus virginiana*) and post oak (*Quercus stellata*) were reduced by experimental intensification of summer drought conditions, but growth of eastern redcedar increased in response to warming, while that of post oak did not (Volder *et al.* 2013). This suggests that encroachment by *Juniperus* spp is likely to accelerate with continued climate change and will further reduce the extent and production of grasslands. Woodland encroachment has already forced the reassessment of fire regimes and implementation of prescribed-burning programs within this region (Twidwell *et al.* 2013).

Northern Great Plains

Warming and increased precipitation, coupled with elevated atmospheric CO₂ concentrations, are expected to increase plant production by alleviating temperature and water limitations on plant growth and extending the growing season (Figure 1; Polley *et al.* 2013). Such an outcome would presumably improve the efficiency of livestock production by reducing the period of winter feeding and allowing for increased stocking rates (CCSP 2008). However, it is uncertain to what extent elevated atmospheric CO₂ will reduce forage quality, and thus livestock production and profitability, by increasing plant C:N ratios. Nitrogen concentrations of live plant tissues less than 1.5% are likely to reduce animal growth and reproduction, while values of approximately 1% will be sufficient to meet maintenance requirements of mature animals. The adverse effects of low nutritive forage can be offset by dietary N supplements, but this will increase both operating costs and labor requirements (Schauer *et al.* 2005).

In addition, these climatic shifts are projected to modify plant community composition by facilitating recruitment and growth of invasive herbaceous plants – including leafy spurge (*Euphorbia esula*) and diffuse knapweed (*Centaurea diffusa*), as well as several species of sub-shrubs – and alter the distributional ranges of other invasive species (Morgan *et al.* 2007). Spotted knapweed (*Centaurea biebersteinii*) may shift toward higher elevations, while leafy spurge may extend its range northward into southern Canada (Bradley



Figure 2. Elevated CO₂ promotes the invasion of Wyoming mixed-grass prairie by Dalmatian toadflax (*Linaria dalmatica*), whereas warming has little effect.

and Wilcove 2009). We anticipate that increased abundance and expanded ranges of exotic invasive species are more likely to adversely affect livestock production than such changes in native species, because exotics are often unpalatable and occasionally toxic to livestock (Figure 2).

Northwest

Warming is projected to decrease soil water availability, especially during late summer, and to reduce plant production in this region (Figure 1; Polley *et al.* 2013). Earlier snowmelt, which now occurs 10–15 days earlier than it did 50 years ago, and reduced stream flow are expected to reduce primary production and modify plant species composition in riparian systems (CCSP 2008). Benefits to livestock production provided by milder, wetter winters may partially offset the negative effects of longer and drier summers, but supplemental feeding may be required if the summer dry period is prolonged. Warming and associated range expansion of invasive annual grasses are likely to produce larger and more frequent fires that function as amplifying feedbacks, promoting further invasion of the exotic annual grasses cheatgrass (*Bromus tectorum*), medusahead (*Taeniatherum caput-medusae*), and red brome (*Bromus madritensis rubens*), especially at higher elevations and in northern portions of their current ranges (Smith *et al.* 2000; Ziska *et al.* 2005). Yellow starthistle (*Centaurea solstitialis*) is expected to become more prolific but to expand its range only marginally (Bradley and Wilcove 2009). The lower palatability and nutrient content of these invasive species will negatively affect livestock production by reducing forage intake and increasing foraging time of livestock (Polley *et al.* 2013).

■ Adaptive capacity

Rangeland managers have historically developed considerable adaptive capacity to contend with economic, environmental, and ecosystem variability. However, climate

Panel 1. An assessment of heterogeneous adaptive capacity among 100 rangeland livestock managers

Adaptive capacity is necessary to convert natural and social resources into useful adaptation strategies, and is considered to include four dimensions: capacity to manage risk and uncertainty; capacity to plan, learn, and reorganize; emotional and financial flexibility to incorporate the costs of change; and interest in adapting to change.

These four dimensions were used to assess the adaptive capacity of 100 livestock managers in northern Australia to address climatic variability (Figure 3; Marshall and Smajgl 2013). In face-to-face interviews, managers were asked a series of questions and were then categorized as possessing low or high capacity for each of the four dimensions. The skill sets of individual managers could include high capacity in all, some, or none of the four dimensions. Variation in adaptive capacity of the 100 managers represented all 16 combinations of the four dimensions.

Managers perceived their ability to address climate risk positively; for example, Marshall and Smajgl (2013) noted that 90% of managers believed that they were “more likely to survive drought compared to other cattle producers”. With respect to capacity to plan, managers appeared to develop plans for risks such as drought based on their own skills and knowledge. Responses varied from “just hope for the best if there is a drought” (21%) to they were “good at doing what [they] do and trust [their] own decision” (90%). Their emotional flexibility was generally greater than their financial flexibility. The managers expressed an interest in climate-change adaptation but no interest in the strategies. While 83% were interested in how they could better prepare for drought, less than 60% were prepared to learn new skills outside of the industry or attend a workshop on how to better manage for drought. Any single initiative to address adaptation to climate change is unlikely to meet the needs of all managers. For instance, some managers could benefit by improving their ability to manage risk, whereas others would benefit more by developing other skills and information networks.




Figure 3. Proportion of ranchers with high (uppercase) or low (lowercase) adaptive capacity as described by four dimensions: (A, a) capacity to manage risk and uncertainty; (B, b) capacity to plan, learn, and reorganize; (C, c) emotional and financial flexibility to incorporate the costs of change; and (D, d) interest in adapting to change. Modified from Marshall and Smajgl (2013).

change is projected to increase the frequency and extremes of climatic fluctuations beyond those previously experienced, which will require even greater adaptive capacity to minimize failure and abandonment of production enterprises. Adaptive capacity – which encompasses the ability to recognize and manage risk, plan and implement adaptation strategies, display financial and emotional flexibility (described in the section below), and even exhibit awareness of climate change and the need for adaptation – has been demonstrated to vary greatly among managers (Panel 1 and Figure 3; Marshall and Smajgl 2013). We reference the Australian survey conducted by Marshall and Smajgl (2013) because it is the most comprehensive assessment of heterogeneous adaptive capacity among livestock managers. The attributes discussed in that survey are often used to characterize adaptive capacity, and thus we consider the survey results to be broadly applicable to US rangelands and rangeland managers as well. The adaptive capacity of managers, and consequently of livestock operations, will establish the foundation upon which adaptation strategies for climate change are conceptualized, evaluated, and implemented (Fazey *et al.* 2010). Yet many important regional adaptations will far exceed the financial and technical capacity of individual operations and may require public investment (Mendelsohn 2000; IPCC 2014).

Risk management

Managing risks associated with change and uncertainty is fundamental to adaptive capacity and adaptation (Marshall and Stokes 2014). Rangeland livestock managers are continually confronted by risks associated with the unpredictability of markets and weather (Torell *et al.* 2010). Drought management planning can help ranchers minimize reactionary approaches to drought that may result in a loss of productivity and financial assets. Evidence of heterogeneous adaptive capacity is seen in the implementation of current drought management planning. For example, of Utah cattle ranchers surveyed after the 1999 to 2004 drought, only 14% were self-described as adequately prepared. However, drought preparedness increased to 29% of ranchers in 2009, but most remained ill-prepared (Coppock 2011). Sixty percent of Wyoming ranchers incorporated some type of drought management planning, and multiple adaptation strategies were adopted with increasing duration of drought (Kachergis *et al.* 2013). Similarly, nearly 50% of the ranchers surveyed in Australia were unprepared to manage risk and uncertainty associated with periodic drought (Panel 1; Marshall and Smajgl 2013).

Planning and learning for enterprise reorganization

The capacity to plan, learn, and reorganize enterprise structure is dependent upon creativity, experiential and scientific knowledge, awareness of opportunities, and the skill to capitalize on those opportunities (Stafford Smith *et al.* 2007; Joyce *et al.* 2013; Marshall and Stokes 2014). Livestock managers recognize that social capital, in the form of learning networks and environmental awareness, is a more important component of climate-change adaptation than technical information and solutions (Marshall 2010). Ranchers who exhibit foresight and have access to conservation information are most likely to participate in conservation programs (Lubell *et al.* 2013). Consequently, livestock managers will require a variety of adaptation strategies to accommodate the heterogeneous adaptive capacity within this agricultural sector (Panel 1 and Figure 3; Marshall and Smajgl 2013).

Production efficiencies, market pressures, and climate change have led to the restructuring and physical relocation of livestock production systems in parts of the US (Joyce *et al.* 2013; Mu *et al.* 2013). Managers in the hotter areas of Texas, for instance, have increased the proportion of Brahman and Brangus (crosses between Brahman and Angus breeds) cattle in their herds (Zhang *et al.* 2013). This strategy of crossing the two cattle breeds combines consumer preference for Angus beef with the heat and insect tolerance of Brahmans, an adaptation that may become more widely implemented as projections of a warmer and drier climate are realized in Texas and the Southwest US (Hoffmann 2010; Mu *et al.* 2013).

Emotional and financial flexibility

Emotional and financial flexibility of livestock managers are strong indicators of their ability to cope with change. Managers with limited social and natural-resource flexibility are especially vulnerable to weather variation (Marshall 2011) and potentially to climate change. Limited emotional flexibility is indicative of a strong attachment to livestock production as a livelihood and a reluctance to search for employment elsewhere or to diversify livelihoods (Marshall and Smajgl 2013). Often, these managers are older, are highly independent, and lack transferable skills. Financial flexibility, as indicated by income and size of production enterprises, has been shown to influence the acceptance and implementation of innovative adaptations. Managers with larger properties are more likely to implement a greater number of drought management adaptations to increase enterprise flexibility and lessen the adverse consequences of drought (Kachergis *et al.* 2013).

Climate-change awareness

Managers' perceptions of climate change will affect their willingness to develop and adopt various adaptations. Suboptimal investments in adaptive capacity, known as

“adaptation deficits”, have been associated with insufficient awareness of climate-change issues, climate-change denial, and distorted perceptions of risk and current preparedness (Williamson *et al.* 2012). Knowledge and perceptions of climate change were influenced by partisan affiliation, political ideology, and gender. Female managers were more likely to hold more scientifically accurate knowledge about climate change than their male counterparts, regardless of political affiliation (Liu *et al.* 2014).

Adaptation strategies

Many adaptation strategies specific to rangeland livestock production – involving conservative stocking rates, robust drought contingency planning, a shift in livestock breeds or species, management of invasive plants and animal parasites, modified operational structure, and geographic relocation of production enterprises – have been identified (Joyce *et al.* 2013). These strategies vary greatly in the extent of modification, timing of implementation, specificity of impact, potential for success, and, as noted previously, the likelihood of adoption as influenced by varying adaptive capacity among managers.

Drought contingency planning for livestock production systems includes low to moderate grazing intensities, maintaining reserve forage supplies, flexible management of herd size and composition, and procedures for timely destocking when necessary (Joyce *et al.* 2013). In areas of highly variable forage production, maintaining grazing flexibility by shifting from cow–calf enterprises to enterprises using yearling cattle may be critical for economic success; this degree of grazing flexibility, however, incurs additional costs and financial risks that may prove challenging for risk-averse managers (Torell *et al.* 2010). A switch to alternative livestock species represents another viable adaptation. As compared with cattle, smaller ruminant livestock – sheep and goats – are more heat tolerant, require less water, can consume a greater diversity of plant species, and may be better suited to future climates, especially in the southern regions (Polley *et al.* 2013). Nevertheless, the availability of market infrastructure and rancher perceptions about and abilities to manage smaller ruminant livestock may necessitate assistance with market development, as well as training and incentives for managers.

Matching adaptation with adaptive capacity and geographic exposure

The sensitivity of a region to climate change will be determined not only by ecological responses but also by the potential adaptive capacity of managers. For example, in the Southwest, the percentage of all agricultural enterprises with sales less than \$10,000 is higher than in the northern Great Plains (NASS 2012a), implying that managers in the former region may have fewer resources to implement novel, and potentially expensive, adaptations. The percentage of principal enterprise managers

who are > 65 years of age is higher across the Southwest and southern Great Plains than in the northern Great Plains and the Northwest (NASS 2012b), suggesting that managers in the former regions may demonstrate greater attachment to place and less willingness to diversify or relocate enterprises. These social characteristics may indicate a lower adaptive capacity, leading to higher rates of enterprise failure. A large number of such failures could irreversibly modify the infrastructure of the regional livestock industry, if the size of regional livestock herds falls below a threshold of economic viability. Recent drought in the southern Great Plains and the consequential closure of cattle feedlots and beef processing plants in the region represent such a case, which will further constrain livestock marketing opportunities for the remaining managers (Johnston 2014).

Adaptations will be required to minimize climate-change impacts and to exploit opportunities associated with more favorable climatic conditions (Table 1). Adaptations to exploit opportunities in the northern Great Plains will involve development of facilities to support greater livestock numbers, and maintenance of sufficient marketing and pricing strategies. The percentage of agricultural enterprises that reported farming as their primary occupation is highest in the northern Great Plains, and average annual enterprise sales in the region are well above \$10 000 (NASS 2012a, c). This suggests that sufficient financial capital is available for managers to capitalize on opportunities associated with greater livestock pro-

duction, but information addressing enterprise expansion and marketing strategies may be required. Novel agricultural opportunities associated with climate change will likely result in competition for land, water, energy, and labor resources among crop and livestock production systems, which will require assessment of land-use trade-offs.

Current federal programs, such as the Environmental Quality Incentives Program (EQIP), offer technical assistance opportunities for various conservation measures to address local and regional environmental concerns (Reimer and Prokopy 2014). However, these programs do not contain a sufficient variety of options to account for the diverse adaptive capacity of managers. The interaction of heterogeneous adaptive capacity among managers and geographically specific ecological consequences of climate change will be particularly challenging for social organizations that offer assistance based on the “average” enterprise structure or “average” adaptive capacity of managers (Figure 4). Multiple approaches will be required to combine scientific and experiential knowledge, develop partnerships for co-production of new management-relevant knowledge, and incentivize the implementation of adaptation strategies that cater to the needs and abilities of diverse managers and associated stakeholder groups (Dilling and Lemos 2011; Bierbaum *et al.* 2013). Some of these strategies are currently in place, but others are required in the form of social organizations capable of providing risk-management programs and financing for technology and equipment to implement capital-intensive adaptations (IPCC 2014).

Table 1. Two distinct categories of adaptation strategies to contend with the consequences of climate change on US rangelands

Minimize adversity	Optimize opportunity
 <p>Southwestern rangelands</p>	 <p>Northern Great Plains rangelands</p>
Increase awareness of climatic variability and potential extremes	Increase awareness of opportunities presented by climate change
Develop management skills and flexibility to cope with climate extremes	Develop infrastructure to support greater livestock numbers
Shift livestock breeds or species to manage heat stress and parasites	Evaluate market availability and price stability
Use novel and invasive plant species for forage, biofuels, and carbon sequestration	Manage potential increases in invasive plants, and animal parasites and diseases
Seek markets for alternative ecosystem services	Assess land-use trade-offs with crops, biofuels, energy extraction, and biodiversity conservation

■ When incremental adaptation fails

Severity of exposure to climate change, enterprise sensitivity, and available adaptive capacity may interact in some instances to produce conditions where incremental adaptation is insufficient to sustain current livelihoods (Joyce *et al.* 2013). In these cases, production enterprises and even entire human–ecological systems may have to be reorganized to create alternative livelihoods based on different combinations of ecosystem services (Kates *et al.* 2012). For example, enterprise managers may find that it is not economical to compensate for climate-induced declines in livestock production resulting from more frequent and intense drought, accelerated woody plant encroachment, and invasion of exotic herbaceous species (Polley *et al.* 2013). Alternative livelihoods to livestock grazing, including ecotourism, watershed management, and alternative energy sources, may be more compatible with the ecological and social condi-

tions of some regions in the future. Development of frameworks and policies that identify and guide implementation of transformational change is a critical but largely overlooked challenge.

Conclusion

Livelihoods linked to livestock grazing in the western US are especially vulnerable to climate change. Vulnerability is a function of exposure to unique intensities and combinations of climate-change drivers, along with the diverse adaptive capacity of managers to contend with these consequences. The geographic specificity of ecological consequences and the heterogeneous adaptive capacity among enterprise managers represent underappreciated and interacting components of climate-change adaptation. Categories of adaptation strategies will be required to minimize the adverse consequences of climate change in the Southwest US and in Texas, and to capitalize on potential climate-induced opportunities in the northern Great Plains of the US. This establishes a robust justification for the development of an array of flexible and cost-effective adaptation strategies to contend with these diverse ecological consequences, while accounting for the varied adaptive capacity of managers to adopt and implement them. Although a few of these strategies are currently in place, others are not, and their development may require some degree of public involvement given the enormity of the challenge. The widespread occurrence of adaptation deficits in the face of current climatic variability foreshadows major challenges that will be encountered in the development and implementation of adaptation strategies that will be needed to cope with increasing climatic variability.

References

- Baker NC and Huang H. 2014. A comparative study of precipitation and evaporation between CMIP3 and CMIP5 climate model ensembles in semiarid regions. *J Climate* **27**: 3731–49.
- Balch JK, Bradley BA, D'Antonio CM, and Gomez-Dans J. 2013. Introduced annual grass increases regional fire activity across the arid western USA (1980–2009). *Glob Change Biol* **19**: 173–83.
- Bierbaum RJ, Smith B, Lee A, *et al.* 2013. A comprehensive review of climate adaptation in the United States: more than before, but less than needed. *Mitig Adapt Strateg Glob Change* **18**: 361–406.
- Bradley BA and Wilcove DS. 2009. When invasive plants disappear: transformative restoration possibilities in the western United States resulting from climate change. *Restor Ecol* **17**: 715–21.
- CCSP (Climate Change Science Program). 2008. Synthesis and




Figure 4. Geographically specific exposure to climate change and heterogeneous adaptive capacity may interact to increase the complexity and diversity of adaptation strategies necessary to maintain livelihoods dependent on livestock grazing. Adaptation strategies must be sufficiently diverse to minimize adverse consequences and optimize opportunities of climate change, and to accommodate managers who possess varied adaptive capacities. Livelihoods will be sustained only if managers are willing to adopt adaptation strategies that are appropriate for specific regions.

- Assessment Product 4.3 (SAP 4.3): the effects of climate change on agriculture, land resources, water resources, and biodiversity in the United States. Washington, DC: CCSP.
- Coppock DL. 2011. Ranching and multiyear droughts in Utah: production impacts, risk perceptions, and changes in preparedness. *Rangeland Ecol Manag* **64**: 607–18.
- Craine JM, Town EG, Joern A, and Hamilton RG. 2009. Consequences of climate variability for the performance of bison in tallgrass prairie. *Glob Change Biol* **15**: 772–79.
- Dilling L and Lemos MC. 2011. Creating usable science: opportunities and constraints for climate knowledge use and their implications for science policy. *Global Environ Chang* **21**: 680–89.
- Fazey I, Gamarra JBP, Fischer J, *et al.* 2010. Adaptation strategies for reducing vulnerability to future environmental change. *Front Ecol Environ* **8**: 414–22.
- Hoffmann I. 2010. Climate change and the characterization, breeding and conservation of animal genetic resources. *Anim Genet* **41**: S32–S46.
- Howden SM, Crimp SJ, and Stokes CJ. 2008. Climate change and Australian livestock systems: impacts, research and policy issues. *Aust J Exp Agr* **48**: 780–88.
- IPCC (Intergovernmental Panel on Climate Change). 2013. The physical science basis: contribution of Working Group I to the Fifth Assessment Report. www.ipcc.ch/report/ar5/wg1/. Viewed 18 Mar 2015.
- IPCC (Intergovernmental Panel on Climate Change). 2014. Impacts, adaptation and vulnerability: contribution of Working Group II to the Fifth Assessment Report. www.ipcc.ch/report/ar5/wg2/. Viewed 18 Mar 2015.
- Johnston T. 2014. Dry age beef. Chicago, IL: Meatingplace. <http://dryagebeef.meatingplace.com/index.php>. Viewed 23 Jan 2015.
- Joyce LA, Briske DD, Brown JR, *et al.* 2013. Climate change and North American rangelands: assessment of mitigation and adaptation strategies. *Rangeland Ecol Manag* **66**: 512–28.

- Kachergis E, Derner JD, Cutts BB, et al. 2013. Increasing flexibility in rangeland management during drought. *Ecosphere* 5: art77.
- Kates RW, Travis WR, and Wilbanks TJ. 2012. Transformational adaptation when incremental adaptations to climate change are insufficient. *P Natl Acad Sci USA* 109: 7156–61.
- Knapp AK, Beier C, Briske DD, et al. 2008. Consequences of more extreme precipitation regimes for terrestrial ecosystems. *BioScience* 58: 811–21.
- Liu Z, Smith Jr WJ, and Safi AS. 2014. Rancher and farmer perceptions of climate change in Nevada, USA. *Climatic Change* 122: 313–27.
- Lubell MN, Cutts BB, Roche LM, et al. 2013. Conservation program participation and adaptive rangeland decision-making. *Rangeland Ecol Manag* 66: 609–20.
- Marshall NA and Smajgl A. 2013. Understanding variability in adaptive capacity on rangelands. *Rangeland Ecol Manag* 66: 88–94.
- Marshall NA and Stokes CJ. 2014. Influencing adaptation processes on the Australian rangelands for social and ecological resilience. *Ecol Soc* 19: 14.
- Marshall NS. 2010. Understanding social resilience to climate variability in primary enterprises and industries. *Global Environ Chang* 20: 36–43.
- Marshall NS. 2011. Assessing resource dependency on the rangelands as a measure of climate sensitivity. *Soc Natur Resour* 24: 1105–15.
- Meehl GA, Covey C, Delworth T, et al. 2007. The WCRP CMIP3 multi-model dataset: a new era in climate change research. *B Am Meteorol Soc* 88: 1383–94.
- Mendelsohn R. 2000. Efficient adaptation to climate change. *Climatic Change* 45: 583–600.
- Morgan JA, Milchunas DG, LeCain DR, et al. 2007. Carbon dioxide enrichment alters plant community structure and accelerates shrub growth in the shortgrass steppe. *P Natl Acad Sci USA* 104: 14724–29.
- Morgan JA, Moiser AR, Milchunas DG, et al. 2004. CO₂ enhances productivity, alters species composition, and reduces digestibility of shortgrass steppe vegetation. *Ecol Appl* 14: 208–19.
- Mu JE, McCarl BA, and Wein AM. 2013. Adaptation to climate change: changes in farmland use and stocking rate in the US. *Mitig Adapt Strateg Glob Change* 18: 713–30.
- NASS (National Agricultural Statistics Service). 2012a. Percent of farms with sales of less than \$10 000: 2012. Washington, DC: USDA NASS. www.agcensus.usda.gov/Publications/2012/Online_Resources/Ag_Atlas_Maps/Economics/Farms_by_Size/12-M009.php. Viewed 11 Jan 2015.
- NASS (National Agricultural Statistics Service). 2012b. Percent of principal farm operators 65 years old and over: 2012. Washington, DC: USDA NASS. www.agcensus.usda.gov/Publications/2012/Online_Resources/Ag_Atlas_Maps/Operators/Characteristics/12-M124.php. Viewed 11 Jan 2015.
- NASS (National Agricultural Statistics Service). 2012c. Percent of principal farm operators reporting primary occupation as farming: 2012. Washington, DC: USDA NASS. www.agcensus.usda.gov/Publications/2012/Online_Resources/Ag_Atlas_Maps/Operators/Principle_Occupation/12-M120.php. Viewed 11 Jan 2015.
- NRC (National Research Council). 2010. Adapting to climate change. Washington, DC: National Academy Press.
- Polley HW, Briske DD, Morgan JA, et al. 2013. Climate change and North American rangelands: evidence, trends, and implications. *Rangeland Ecol Manag* 66: 493–511.
- Reimer A and Prokopy L. 2014. One federal policy, four different policy contexts: an examination of agri-environmental policy implementation in the midwestern United States. *Land Use Policy* 38: 605–14.
- Reynolds JF, Stafford Smith DM, Lambin EF, et al. 2007. Global desertification: building a science for drylands. *Science* 316: 847–51.
- Sayre NF, McAllister RRR, Bestelmeyer BT, et al. 2013. Earth Stewardship of rangelands: coping with ecological, economic, and political marginality. *Front Ecol Environ* 11: 348–54.
- Schauer CS, Bohnert DW, Ganskopp DC, et al. 2005. Influence of protein supplementation frequency on cows consuming low-quality forage: performance, grazing behavior, and variation in supplement intake. *J Anim Sci* 83: 1715–25.
- Smith SD, Huxman TE, Zitzer SF, et al. 2000. Elevated CO₂ increases productivity and invasive species success in an arid ecosystem. *Nature* 408: 79–82.
- Stafford Smith DM, McKeon GM, Watson IW, et al. 2007. Learning from episodes of degradation and recovery in variable Australian rangelands. *P Natl Acad Sci USA* 104: 20690–95.
- Taylor KE, Stouffer RJ, and Meehl GA. 2012. An overview of CMIP5 and the experiment design. *B Am Meteorol Soc* 93: 485–98.
- Torell LA, Murugan S, and Ramirez OA. 2010. Economics of flexible versus conservative stocking strategies to manage climate variability risk. *Rangeland Ecol Manag* 63: 415–25.
- Twidwell D, Rogers WE, Fuhlendorf SD, et al. 2013. The rising Great Plains fire campaign: citizens' response to woody plant encroachment. *Front Ecol Environ* 11: e64–e71; doi:10.1890/130015.
- USGCRP (US Global Change Research Project). 2009. Global climate change impacts in the United States. In: Karl TR, Melillo JM, and Peterson TC (Eds). Cambridge, UK: Cambridge University Press.
- Volder A, Briske DD, and Tjoelker MG. 2013. Climate warming and precipitation redistribution modify tree–grass interactions and tree species establishment in a warm-temperate savanna. *Glob Change Biol* 19: 843–57.
- Walther GR. 2003. Plants in a warmer world. *Perspect Plant Ecol* 6/3: 169–85.
- Williamson T, Hessel H, and Johnston M. 2012. Adaptive capacity deficits and adaptive capacity of economic systems in climate change vulnerability assessment. *Forest Policy Econ* 15: 160–66.
- Zhang YW, Hagerman AD, and McCarl BA. 2013. Influence of climate factors on spatial distribution of Texas cattle breeds. *Climatic Change* 118: 183–95.
- Ziska LH, Reeves JB, and Blank B. 2005. The impact of recent increases in atmospheric CO₂ on biomass production and vegetative retention of cheatgrass (*Bromus tectorum*): implications for fire disturbance. *Glob Change Biol* 11: 1325–32.

⁴US Department of Agriculture, Jornada Experimental Range, Natural Resource Conservation Service, New Mexico State University, Las Cruces, NM; ⁵University of Colorado and National Oceanic and Atmospheric Administration, Earth Systems Research Laboratory, Boulder, CO; ⁶US Department of Agriculture–Agricultural Research Service, Crops Research Laboratory, Fort Collins, CO; ⁷Department of Agricultural Economics, Texas A&M University, College Station, TX; ⁸Animal and Range Sciences Department, New Mexico State University, Las Cruces, NM